

When a partial Borel order is linearizable

Vladimir Kanovei ^{*†‡}

April 1997

Abstract

We prove the following classification theorem of the “Glimm – Effros” type for Borel order relations: a Borel partial order on the reals either is Borel linearizable or includes a copy of a certain Borel partial order $\leq_0$ which is not Borel linearizable.

^{*} Moscow Transport Engineering Institute

[†] `kanovei@mech.math.msu.su` and `kanovei@math.uni-wuppertal.de`

[‡] This paper was accomplished during my visit to Caltech in April 1997. I thank Caltech for the support and A. S. Kechris and J. Zapletal for useful information and interesting discussions relevant to the topic of this paper during the visit.

Notation

A binary relation $\preceq$ on a set X is a *partial quasi-order*, or *p. q.-o.* in brief, on X , iff $x \preceq y \wedge y \preceq z \implies x \preceq z$, and $x \preceq x$ for any $x \in X$. In this case, $\approx$ is the associated equivalence relation, *i. e.* $x \approx y$ iff $x \preceq y \wedge y \preceq x$.

If in addition $x \approx x \implies x = x$ for any x then $\preceq$ is a *partial order*, or *p. o.*, so that, say, forcing relations are p. q.-o.'s, but, generally speaking, not p. o.'s in this notational system.

A p. o. is *linear* (l. o.) iff we have $x \preceq y \vee y \preceq x$ for all $x, y \in X$.

Let $\preceq$ and $\preceq'$ be p. q.-o.'s on resp. X and X' . A map $h : X \longrightarrow X'$ will be called *half order preserving*, or *h. o. p.*, iff $x \preceq y \implies h(x) \preceq' h(y)$.

Definition 1 A Borel p. q.-o. $\langle X; \preceq \rangle$ is *Borel linearizable* iff there is a Borel l. o. $\langle X'; \preceq' \rangle$ and a Borel h. o. p. map $h : X \longrightarrow X'$ (called: *linearization map*) satisfying $x \approx y \iff h(x) = h(y)$.¹

Introduction

Harrington, Marker, and Shelah [2] proved several theorems on Borel partial orders, mainly concerning *thin* p. q.-o.'s, *i. e.* those which do not admit uncountable pairwise incomparable subsets. In particular they demonstrated that any such a Borel p. q.-o. is Borel linearizable, and moreover the corresponding l. o. $\langle X'; \preceq' \rangle$ can be chosen so that $X' \subseteq 2^\alpha$ for some $\alpha < \omega_1$ while $\preceq' = \leq_{\text{lex}} \upharpoonright X$, where $\leq_{\text{lex}}$ is the lexicographical order.

As elementary examples show that the thinness is not a necessary condition for the Borel linearization, this result leaves open the problem of linearization of non-thin Borel p. o.'s. Harrington *e. a.* write in [2] that “there is little to say about nonthin orderings”, although there are many interesting among them like the *dominance* order on ω^ω .

Our main result will say that not all Borel p. q.-o.'s are Borel linearizable, and there exists a *minimal* in certain sense among them.

Definition 2 Let $a, b \in 2^\omega$. We define $a \leq_0 b$ iff either $a = b$ or $a \mathbf{E}_0 b$ ² and $a(k_0) < b(k_0)$ where k_0 is the largest k such that $a(k) \neq b(k)$.³

The relation $\leq_0$ is a Borel p. q.-o. on 2^ω which orders every $\mathbf{E}_0$ -class similarly to the integers $\mathbb{Z}$ (except for the class $[\omega \times \{0\}]_{\mathbf{E}_0}$ ordered as ω and the class $[\omega \times \{1\}]_{\mathbf{E}_0}$ ordered as ω^*) but leaves any two $\mathbf{E}_0$ -inequivalent reals incomparable.

¹ The equivalence cannot be dropped as otherwise a one-element set X' works in any case.

² That is $a(k) = b(k)$ for all but finite k — the *Vitali* equivalence relation on 2^ω .

³ If one enlarges $<_0$ so that, in addition, $a <_0 b$ whenever $a, b \in 2^\omega$ are such that $a(k) = 1$ and $b(k) = 0$ for all but finite k then the enlarged relation can be induced by a Borel action of $\mathbb{Z}$ on 2^ω , such that $a <_0 b$ iff $a = zb$ for some $z \in \mathbb{Z}$, $z > 0$.

Theorem 3 (The main result.) *Suppose that $\preceq$ is a Borel p. q.-o. on $\mathcal{N} = \omega^\omega$. Then exactly one of the following two conditions is satisfied:*

- (I) *$\preceq$ is Borel linearizable, moreover there exist an ordinal $\alpha < \omega_1$ and a Borel linearization map $h : \langle \mathcal{N}; \preceq \rangle \longrightarrow \langle 2^\alpha; \leq_{\text{lex}} \rangle$.*
- (II) *there exists a continuous 1 – 1 map $F : 2^\omega \longrightarrow \mathcal{N}$ such that we have $a \leq_0 b \implies F(a) \preceq F(b)$ while $a \not\leq_0 b$ implies that $F(a)$ and $F(b)$ are $\preceq$ -incomparable.*⁴

The theorem resembles the case of Borel equivalence relations where a necessary and sufficient condition for a Borel equivalence relation E to be *smooth* is that E_0 (which is not smooth) does not continuously embed in E (Harrington, Kechris, Louveau [1]). ($\leq_0$ itself is *not* Borel linearizable.)

The proof is essentially a combination of ideas and technique in [1, 2].

1 Incompatibility

Let us first prove that (I) and (II) are incompatible.

Suppose otherwise. The superposition of the maps F and h is then a Borel h. o. p. map $\phi : \langle 2^\omega; \leq_0 \rangle \longrightarrow \langle 2^\alpha; \leq_{\text{lex}} \rangle$ satisfying the following: $\phi(a) = \phi(b)$ implies that $a E_0 b$, i. e. a and b are $\leq_0$ -comparable.

Therefore, as any E_0 -class is $\leq_0$ -ordered similarly to $\mathbb{Z}$, ω , or ω^* , the ϕ -image $X_a = \phi'' [a]_{E_0}$ of the E_0 -class of any $a \in 2^\omega$ is $\leq_{\text{lex}}$ -ordered similarly to a subset of $\mathbb{Z}$. If $X_a = \{x_a\}$ is a singleton then put $\psi(a) = x_a$.

Assume now that X_a contains at least two points. In this case we can effectively pick an element in X_a ! Indeed there is a maximal sequence $u \in 2^{<\alpha}$ such that $u \subset x$ for each $x \in X_a$. Then the set $X_a^{\text{left}} = \{x \in X : u \wedge 0 \subset x\}$ contains a $\leq_{\text{lex}}$ -largest element, which we denote by $\psi(a)$.

To conclude ψ is a Borel reduction of E_0 to the equality on 2^α , i. e. $a E_0 b$ iff $\psi(a) = \psi(b)$, which is impossible.

2 The dichotomy

As usual it will be assumed that the p. q.-o. $\preceq$ of Theorem 3 is a Δ_1^1 relation. Let $\approx$ denote the associated equivalence.

Following [2] let, for $\alpha < \omega_1^{\text{CK}}$, $\mathcal{F}_\alpha$ to be the family of all h. o. p. Δ_1^1 functions $f : \langle \mathcal{N}; \preceq \rangle \longrightarrow \langle 2^\alpha; \leq_{\text{lex}} \rangle$. Then $\mathcal{F} = \bigcup_{\alpha < \omega_1^{\text{CK}}} \mathcal{F}_\alpha$ is a (countable) Π_1^1 set, in a suitable coding system for functions of this type. (See [2] on details.)

⁴ Then F associates a chain $\{F(b) : b E_0 a\}$ in $\langle \mathcal{N}; \preceq \rangle$ to each E_0 -class $[a]_{E_0}$ so that any two different chains do not contain $\preceq$ -comparable elements: let us call them *fully incomparable* chains. Thus (II) essentially says that $\preceq$ admits an effectively “big” Borel family of fully incomparable chains, which is therefore necessary and sufficient for $\preceq$ to be *not* Borel linearizable.

Define, for $x, y \in \mathcal{N}$, $x \equiv y$ iff $f(x) = f(y)$ for any $f \in \mathcal{F}$.

Lemma 4 (See [2].) $\equiv$ is a Σ_1^1 equivalence relation including $\approx$.

Proof As $\preceq$ is Δ_1^1 , one gets by a rather standard argument a set $N \subseteq \omega$ and a function $f_n \in \mathcal{F}$ for any $n \in N$ so that $\mathcal{F} = \{f_n : n \in N\}$ and the relations $n \in N \wedge f_n(x) \leq_{\text{lex}} f_n(y)$ and $n \in N \wedge f_n(x) <_{\text{lex}} f_n(y)$ are presentable in the form $n \in N \wedge \mathcal{O}(x, y)$ and $n \in N \wedge \mathcal{O}'(x, y)$ where $\mathcal{O}, \mathcal{O}'$ are Σ_1^1 relations. Now $x \equiv y$ iff $\forall n (n \in N \implies f_n(x) = f_n(y))$, as required. $\square$

Case 1: $\equiv$ coincides with $\approx$.

Let us show how this implies (I) of Theorem 3. The set

$$P = \{\langle x, y, n \rangle : x \not\approx y \wedge f_n(x) \neq f_n(y)\}.$$

is Π_1^1 and, by the assumption of Case 1, its projection on x, y coincides with the complement of $\approx$. Let $Q \subseteq P$ be a Π_1^1 set uniformizing P in the sense $\mathcal{N}^2 \times \omega$. Then Q is Δ_1^1 because $Q(x, y, n) \iff x \not\approx y \wedge \forall n' \neq n \neg Q(x, y, n')$. It follows that $N' = \{n : \exists x, y Q(x, y, n)\} \subseteq N$ is Σ_1^1 . Therefore there is a Δ_1^1 set M such that $N' \subseteq M \subseteq N$.⁵

Consider a Δ_1^1 enumeration $M = \{n_l : l \in \omega\}$. For any l , $f_{n_l} \in \mathcal{F}_\alpha$ for some ordinal $\alpha = \alpha_l < \omega_1^{\text{CK}}$. Another standard argument (see [2]) shows that in this case (*e.g.* when $M \subseteq N$ is Δ_1^1) the ordinals α_l are bounded by some $\alpha < \omega_1^{\text{CK}}$. It follows that the function $h(x) = f_{n_0}(x) \wedge f_{n_1}(x) \wedge f_{n_2}(x) \wedge \dots \wedge f_{n_l}(x) \wedge \dots$ belongs to some $\mathcal{F}_\beta$, $\beta \leq \alpha \cdot \omega$. On the other hand, by the construction we have $x \approx y \iff h(x) = h(y)$ so h satisfies (I) of Theorem 3.

Case 2: $\approx \subsetneq \equiv$. Assuming this we work towards (II) of Theorem 3.

3 The domain of singularity

By the assumption the Σ_1^1 set $A = \{x : \exists y (x \approx y \wedge x \not\equiv y)\}$ is non-empty.

Define $X \equiv Y$ iff $\forall x \in X \exists y \in Y x \equiv y$ and *vice versa*.

Proposition 5 Let $X, Y \subseteq A$ be non-empty Σ_1^1 sets satisfying $X \equiv Y$. Then

$$P_+ = \{\langle x, y \rangle \in X \times Y : x \equiv y \wedge x \preceq y\}, \quad P_- = \{\langle x, y \rangle \in X \times Y : x \equiv y \wedge x \not\preceq y\}$$

are non-empty Σ_1^1 sets, their projections $\text{pr}_1 P_+$ and $\text{pr}_1 P_-$ are Σ_1^1 -dense in X ⁶ while the projections $\text{pr}_2 P_+$ and $\text{pr}_2 P_-$ are Σ_1^1 -dense in Y .⁷

⁵ Harrington *e. a.* [2] use a general reflection theorem to get such a set, but a more elementary reasoning sometimes has advantage.

⁶ That is intersect any non-empty Σ_1^1 set $X' \subseteq X$.

⁷ For a set $P \subseteq \mathcal{N}^2$, $\text{pr}_1 P$ and $\text{pr}_2 P$ have the obvious meaning of the projections on the resp. 1st and 2nd copy of $\mathcal{N}$.

Proof The density easily follows from the non-emptiness, so let us concentrate on the latter. *Prove that $P_+ \neq \emptyset$.*

Suppose on the contrary that $P_+ = \emptyset$. Then there is a single function $f \in \mathcal{F}$ such that the set $\{\langle x, y \rangle \in X \times Y : f(x) = f(y) \wedge x \preceq y\}$ is empty. (See the reasoning in Case 1 of Section 2.) Define

$$X_\infty = \{x : \forall y \in Y (f(x) = f(y) \implies x \not\preceq y)\},$$

so that X_∞ is a Π_1^1 set and $X \subseteq X_\infty$ but $Y \cap X_\infty = \emptyset$. Using Separation we can easily define a sequence of sets

$$X = X_0 \subseteq U_0 \subseteq X_1 \subseteq U_1 \subseteq \dots \subseteq X_n \subseteq U_n \subseteq \dots \subseteq X_\infty$$

so that $U_n = \{x' : \exists x \in X_n (f(x) = f(x') \wedge x \preceq x')\}$ while $X_{n+1} \in \Delta_1^1$ for all n . (Note that if $X_n \subseteq X_\infty$ and U_n is defined as indicated then $U_n \subseteq X_\infty$ too.) Moreover a proper execution of the construction⁸ allows to get the final set $U = \bigcup_n U_n = \bigcup_n X_n$ in Δ_1^1 . Note that $X \subseteq U$ but $Y \cap U = \emptyset$ as $U \subseteq X_\infty$.

Now put $f'(x) = f(x) \wedge 1$ whenever $x \in U$ and $f'(x) = f(x) \wedge 0$ otherwise. *We assert that $f' \in \mathcal{F}$.* Indeed suppose that $x' \preceq y'$ and prove $f'(x') \leq_{\text{lex}} f'(y')$. It can be assumed that $f(x') = f(y')$. It remains to check $x' \in U \implies y' \in U$, which easily follows from the definition of sets U_n . Thus $f' \in \mathcal{F}$.

However clearly $f'(x) \neq f'(y)$, hence $x \not\equiv y$, whenever $x \in X$ and $y \in Y$ which is a contradiction with the assumption that $X \equiv Y$.

Now *prove that $P_- \neq \emptyset$.* Consider first the case $X = Y$. Suppose on the contrary that $P_- = \emptyset$. Then, as above, there is a single function $f \in \mathcal{F}$ such that the set $\{\langle x, y \rangle \in X^2 : f(x) = f(y) \wedge x \not\preceq y\}$ is empty, so that $\equiv$ and $\approx$ coincide on X . Our plan is to find functions $f', f'' \in \mathcal{F}$ such that the sets

$$\begin{aligned} Q' &= \{\langle x, y \rangle \in X \times \mathcal{N} : f'(x) = f'(y) \wedge y \not\preceq x\} \\ Q'' &= \{\langle x, y \rangle \in X \times \mathcal{N} : f''(x) = f''(y) \wedge x \not\preceq y\} \end{aligned}$$

are empty; then $Q = \{\langle x, y \rangle \in X \times \mathcal{N} : x \equiv y \wedge y \not\approx x\} = \emptyset$, which a contradiction with $\emptyset \neq X \subseteq A$.

Let us find f' ; the other case is similar. Define

$$X_\infty = \{x : \forall x' \in X (f(x) = f(x') \implies x \preceq x')\},$$

so that X_∞ is Π_1^1 and $X \subseteq X_\infty$. As above there is a sequence of sets

$$X = X_0 \subseteq U_0 \subseteq X_1 \subseteq U_1 \subseteq \dots \subseteq X_n \subseteq U_n \subseteq \dots \subseteq X_\infty$$

⁸ We refer to the proof of an “invariant” effective Separation theorem in Harrington, Kechris, Louveau [1], which includes a similar construction.

such that $U_n = \{u : \exists x \in X_n (f(x) = f(u) \wedge u \preceq x)\}$ while $X_{n+1} \in \Delta_1^1$ for all n and the final set $U = \bigcup_n U_n = \bigcup_n X_n$ belongs to Δ_1^1 .

Set $f'(x) = f(x) \wedge 0$ whenever $x \in U$ and $f'(x) = f(x) \wedge 1$ otherwise. Then $f' \in \mathcal{F}$. Prove that f' witnesses that $Q' = \emptyset$. Consider any $x \in X$ and $y \in \mathcal{N}$ such that $f'(x) = f'(y)$. Then in particular $f(x) = f(y)$ and $x \in U \iff y \in U$, so that $y \in U$ because we know that $x \in X \subseteq U$. Thus $U \in X_\infty$, so by definition $y \preceq x$ as required.

Finally prove $P_- \neq \emptyset$ in the general case. By the result in the particular case, the Σ_1^1 set $P' = \{\langle x, x' \rangle \in X^2 : x \equiv x' \wedge x \not\preceq x'\}$ is non-empty. Let $X' = \{x' \in X : \exists x P'(x, x')\}$ and $Y' = \{y \in Y : \exists x' \in X' (x' \equiv y)\}$, so that X' and Y' are Σ_1^1 sets satisfying $X' \equiv Y'$. By the result for P_+ there exist $x' \in X'$ and $y \in Y'$ satisfying $x' \equiv y$ and $y \preceq x'$. Now there exists $x \in X$ such that $x \equiv x'$ and $x \not\preceq x'$. Then $x \equiv y$ and $x \not\preceq y$, as required. $\square$

4 The forcing notions involved

Our further strategy will be the following. We shall define a generic extension of the universe $\mathbf{V}$ (where Theorem 3 is being proved) in which there exists a function F which witnesses (II) of Theorem 3. However as the existence of such a function is a Σ_2^1 statement, we obtain the result for $\mathbf{V}$ by Shoenfield.⁹

Definition 6 $\mathbb{P}$ is the collection of all non-empty Σ_1^1 sets $X \subseteq A$.

It is a standard fact that $\mathbb{P}$ (the *Gandy forcing*) forces a real which is the only real which belongs to every set in the generic set $G \subseteq \mathbb{P}$. (We identify Σ_1^1 sets in the universe $\mathbf{V}$ with their copies in the extension.)

Definition 7 $\mathbb{P}_2^+$ is the collection of all non-empty Σ_1^1 sets $P \subseteq A^2$ such that $P(x, y) \implies x \equiv y \wedge x \preceq y$. $\mathbb{P}_2^-$ is defined similarly but with $x \not\preceq y$ instead.

Both of them force a pair of reals $\langle x, y \rangle \in A^2$ satisfying resp. $x \preceq y$ and $x \not\preceq y$. ($\mathbb{P}_2^+$ and $\mathbb{P}_2^-$ are non-empty forcing notions by Proposition 5.)

Definition 8 $\mathbb{P}_\equiv^2$ is the collection of all sets of the form $\Upsilon = X \times Y$ where X, Y are sets in $\mathbb{P}$ satisfying $X \equiv Y$.

Lemma 9 $\mathbb{P}_\equiv^2$ forces a pair of reals $\langle x, y \rangle$ such that $x \not\preceq y$.

⁹ In fact the proof can be conducted without any use of metamathematics, as in Harrington *e. a.* [1], but at the cost of longer reasoning.

Proof Suppose that on the contrary a condition $\Upsilon_0 = X_0 \times Y_0$ in $\mathbb{P}_{\equiv}^2$ forces $x \preceq y$. Consider a more complicated forcing $\mathfrak{P}$ which consists of forcing conditions of the form $\mathbf{p} = \langle \Upsilon, P, \Upsilon', Q \rangle$, where $\Upsilon = X \times Y$ and $\Upsilon' = X' \times Y'$ belong to $\mathbb{P}_{\equiv}^2$, $P \in \mathbb{P}_2^+$, $P \subseteq Y \times X'$, $Q \in \mathbb{P}_2^-$, $Q \subseteq X \times Y'$, and the sets $\text{pr}_1 P \subseteq Y$, $\text{pr}_2 P \subseteq X'$, $\text{pr}_1 Q \subseteq X$ and $\text{pr}_2 Q \subseteq Y'$ are Σ_1^1 -dense in resp. Y , X' , X , Y' .

For instance setting $P_0 = \{\langle y, x' \rangle \in Y_0 \times X_0 : y \equiv x' \wedge y \preceq x'\}$ and $Q_0 = \{\langle x, y' \rangle \in X_0 \times Y_0 : x \equiv y' \wedge x \not\preceq y'\}$ we get a condition $\mathbf{p}_0 = \langle \Upsilon_0, P_0, \Upsilon_0, Q_0 \rangle \in \mathfrak{P}$ by Proposition 5.

It is the principal fact that if $\mathbf{p} = \langle \Upsilon, P, \Upsilon', Q \rangle \in \mathfrak{P}$ and we strengthen one of the components within the corresponding forcing notion then this can be appropriately reflected in the other components. To be concrete assume that, for instance, $P^* \in \mathbb{P}_2^+$, $P^* \subseteq P$, and find a condition $\mathbf{p}_1 = \langle \Upsilon_1, P_1, \Upsilon'_1, Q_1 \rangle \in \mathfrak{P}$ satisfying $\Upsilon_1 \subseteq \Upsilon$, $\Upsilon'_1 \subseteq \Upsilon'$, $P_1 \subseteq P^*$, and $Q_1 \subseteq Q$.

Assume that $\Upsilon = X \times Y$ and $\Upsilon' = X' \times Y'$. Consider the non-empty Σ_1^1 sets $Y_2 = \text{pr}_1 P^* \subseteq Y$ and $X_2 = \{x \in X : \exists y \in Y_2 x \equiv y\}$. It follows from Proposition 5 that $Q_1 = \{\langle x, y \rangle \in Q : x \in X_2\} \neq \emptyset$, hence Q_1 is a condition in $\mathbb{P}_2^-$ and $X_1 = \text{pr}_1 Q_1$ is a non-empty Σ_1^1 subset of $X_2 \subseteq X$.

The set $Y_1 = \{y \in Y_2 : \exists x \in X_1 x \equiv y\}$ satisfies $X_1 \equiv Y_1$, therefore $\Upsilon_1 = X_1 \times Y_1 \in \mathbb{P}_{\equiv}^2$. Furthermore $P_1 = \{\langle y, x \rangle \in P^* : y \in Y_1\} \in \mathbb{P}_2^+$.

Put $X'_1 = \text{pr}_2 P_1 \subseteq X'$ and $Y'_1 = \text{pr}_2 Q_1 \subseteq Y'$. Take notice that $Y_1 \equiv X'_1$ because any condition in $\mathbb{P}_2^+$ is a subset of $\equiv$, similarly $X_1 \equiv Y'_1$, and $X_1 \equiv Y_1$, see above. It follows that $X'_1 \equiv Y'_1$, hence $\Upsilon'_1 = X'_1 \times Y'_1$ is a condition in $\mathbb{P}_{\equiv}^2$.

Now $\mathbf{p}_1 = \langle \Upsilon_1, P_1, \Upsilon'_1, Q_1 \rangle \in \mathfrak{P}$ as required.

We conclude that $\mathfrak{P}$ forces “quadruples” of reals $\langle x, y, x', y' \rangle$ such that the pairs $\langle x, y \rangle$ and $\langle x', y' \rangle$ are $\mathbb{P}_{\equiv}^2$ -generic, hence satisfy $x \preceq y$ and $x' \preceq y'$ provided the generic set contains Υ_0 — by the assumption above. Furthermore the pair $\langle y, x' \rangle$ is $\mathbb{P}_2^+$ -generic, hence $y \preceq x'$, while the pair $\langle x, y' \rangle$ is $\mathbb{P}_2^-$ -generic, hence $x \not\preceq y'$, which is a contradiction. $\square$

5 The splitting construction

Let, in the universe $\mathbf{V}$, $\kappa = \mathfrak{c}$. Let $\mathbf{V}^+$ be a κ -collapse extension of $\mathbf{V}$.

Our aim is to define, in $\mathbf{V}^+$, a splitting system of sets which leads to a function F satisfying (II) of Theorem 3. Let us fix two points before the construction starts.

First, as the forcing notions involved are countable in $\mathbf{V}$, there exist, in $\mathbf{V}^+$, enumerations $\{D(n) : n \in \omega\}$, $\{D_2(n) : n \in \omega\}$, and $\{D^2(n) : n \in \omega\}$, of all open dense sets in resp. $\mathbb{P}$, $\mathbb{P}_2^+$, $\mathbb{P}_{\equiv}^2$, which (the dense sets) belong to $\mathbf{V}$, such that $D(n+1) \subseteq D(n)$ etc. for each n .

Second, we introduce the notion of a crucial pair. A pair $\langle u, v \rangle$ of binary sequences $u, v \in 2^n$ is called *crucial* iff $u = 1^k \wedge 0^{\wedge} w$ and $v = 0^k \wedge 1^{\wedge} w$ for some $k < n$ and $w \in 2^{n-k-1}$. One easily sees that the graph of all crucial pairs in 2^n is actually a chain connecting all members of 2^n .

We define, in $\mathbf{V}^+$, a system of sets $X_u \in \mathbb{P}$, where $u \in 2^{<\omega}$, and sets $P_{uv} \in \mathbb{P}_2^+$, $\langle u, v \rangle$ being a crucial pair in some 2^n , satisfying the following conditions:

- (1) $X_u \in D(n)$ whenever $u \in 2^n$; $X_{u \wedge i} \subseteq X_u$;
- (2) if $\langle u, v \rangle$ is a crucial pair in 2^n then $P_{uv} \in D_2(n)$; $P_{u \wedge i, v \wedge i} \subseteq P_{uv}$;
- (3) if $u, v \in 2^n$ and $u(n-1) \neq v(n-1)$ then $X_u \times X_v \in \mathbb{P}_{\equiv}^2$, moreover $\in D^2(n)$ and $X_u \cap X_v = \emptyset$;
- (4) if $\langle u, v \rangle$ is a crucial pair in 2^n then $\text{pr}_1 P_{uv} = X_u$ and $\text{pr}_2 P_{uv} = X_v$.

Why this implies the existence of a required function ?

First of all for any $a \in 2^\omega$ (in $\mathbf{V}^+$) the sequence of sets $X_{a \upharpoonright n}$ is $\mathbb{P}$ -generic over $\mathbf{V}$ by (1), therefore the intersection $\bigcap_{n \in \omega} X_{a \upharpoonright n}$ is a singleton. Let $F(a) \in \mathcal{N}$ be its only element.

It does not take much effort to prove that F is continuous and $1-1$.

Consider $a, b \in 2^\omega$ satisfying $a \not\leq_0 b$. Then $a(n) \neq b(n)$ for infinitely many n , hence the pair $\langle F(a), F(b) \rangle$ is $\mathbb{P}_{\equiv}^2$ -generic by (3), thus $F(a)$ and $F(b)$ are $\preceq$ -incomparable by Lemma 9.

Consider $a, b \in 2^\omega$ satisfying $a \leq_0 b$. We may assume that a and b are $\leq_0$ -neighbours, *i.e.* $a = 1^k \wedge 0 \wedge c$ while $b = 0^k \wedge 1 \wedge c$ for some $k \in \omega$ and $c \in 2^\omega$. Then by (2) the sequence of sets $P_{a \upharpoonright n, b \upharpoonright n}$, $n > k$, is $\mathbb{P}_2^+$ -generic, hence it results in a pair of reals satisfying $x \preceq y$. However $x = F(a)$ and $y = F(b)$ by (4).

The construction of a splitting system

We argue in $\mathbf{V}^+$.

Suppose that the construction has been completed up to a level n , and expand it to the next level. From now on s, t will denote sequences in 2^n while u, v will denote sequences in 2^{n+1} .

To start with we set $X_{s \wedge i} = X_s$ for all $s \in 2^n$ and $i = 0, 1$, and $P_{s \wedge i, t \wedge i} = P_{st}$ whenever $i = 0, 1$ and $\langle s, t \rangle$ is a crucial pair in 2^n .

For the “initial” crucial pair $\langle 1^n \wedge 0, 0^n \wedge 1 \rangle$ at this level let $P_{1^n \wedge 0, 0^n \wedge 1} = X_{1^n \wedge 0} \times X_{0^n \wedge 1} = X_{1^n} \times X_{0^n}$. Then $P_{1^n \wedge 0, 0^n \wedge 1} \in \mathbb{P}_{\equiv}^2$.¹⁰

This ends the definition of “initial values” at the $n+1$ -th level. The plan is to gradually “shrink” the sets in order to fulfill the requirements.

Step 1. We take care of item (1). Consider an arbitrary $u_0 = s_0 \wedge i \in 2^{n+1}$. As $D(n)$ is dense there is a set $X' \in D(n)$, $X' \subseteq X_{u_0}$. The intension is to take X' as

¹⁰ It easily follows from (2) and (4) that $X_s \equiv X_t$ for all $s, t \in 2^n$, because s and t are connected in 2^n by a unique chain of crucial pairs.

the “new” X_{u_0} . But this change has to be expanded through the chain of crucial pairs, in order to preserve (4).

Thus put $X'_{u_0} = X'$. Suppose that X'_u has been defined and is included in X_u , the old version, for some $u \in 2^{n+1}$, and $\langle u, v \rangle$ is a crucial pair, $v \in 2^{n+1}$ being not yet encountered. Define $P'_{uv} = (X'_u \times \mathcal{N}) \cap P_{uv}$ and $X'_v = \text{pr}_2 P'_{uv}$. Clearly (4) holds for the new sets X'_u , X'_v , and P'_{uv} .

The construction describes how the original change from X_{u_0} to X'_{u_0} spreads through the chain of crucial pairs in 2^{n+1} , resulting in a system of new sets, X'_u and P'_{uv} , which satisfy (1) for the particular $u_0 \in 2^{n+1}$. We iterate this construction consecutively for all $u_0 \in 2^{n+1}$, getting finally a system of sets satisfying (1) (fully) (and (4)), which we shall denote by X_u and P_{uv} from now on.

Step 2. We take care of item (3). Fix a pair of u_0 and v_0 in 2^{n+1} , such that $u_0(n) = 0$ and $v_0(n) = 1$. By the density of $D^2(n)$, there is a set $X'_{u_0} \times X'_{v_0} \in D^2(n)$ included in $X_{u_0} \times X_{v_0}$. We may assume that $X'_{u_0} \cap X'_{v_0} = \emptyset$. (Indeed it easily follows from Proposition 5, for P_- , that there exist reals $x_0 \in X_{u_0}$ and $y_0 \in X_{v_0}$ satisfying $x_0 \equiv y_0$ but $x_0 \neq y_0$, say $x_0(k) = 0$ while $y_0(k) = 1$. Define

$$X = \{x \in X_0 : x(k) = 0 \wedge \exists y \in Y_0 (y(k) = 1 \wedge x \equiv y)\},$$

and Y correspondingly; then $X \equiv Y$ and $X \cap Y = \emptyset$.)

Spread the change from X_{u_0} to X'_{u_0} and from X_{v_0} to X'_{v_0} through the chain of crucial pairs in 2^{n+1} , by the method of Step 1, until the wave of spreading from u_0 meets the wave of spreading from v_0 at the “meeting” crucial pair $\langle 1^n \wedge 0, 0^n \wedge 1 \rangle$. This leads to a system of sets X'_u and P'_{uv} which satisfy (3) for the particular pair $\langle u_0, v_0 \rangle$ and still satisfy (4) possibly except for the “meeting” crucial pair $\langle 1^n \wedge 0, 0^n \wedge 1 \rangle$ (for which basically $P'_{1^n \wedge 0, 0^n \wedge 1}$ is not yet defined for this step).

Note that Step 1 leaves $P_{1^n \wedge 0, 0^n \wedge 1}$ in the form $X_{1^n \wedge 0} \times X_{0^n \wedge 1}$ (where $X_{1^n \wedge 0}$ and $X_{0^n \wedge 1}$ are the “versions” at the end of Step 1). We now have new sets, $X'_{1^n \wedge 0}$ and $X'_{0^n \wedge 1}$, included in resp. $X_{1^n \wedge 0}$ and $X_{0^n \wedge 1}$ and satisfying $X'_{0^n \wedge 0} \equiv X'_{0^n \wedge 1}$ (because we had $X'_{u_0} \equiv X'_{v_0}$ at the beginning of the change.) It remains to define $P'_{1^n \wedge 0, 0^n \wedge 1} = X'_{1^n \wedge 0} \times X'_{0^n \wedge 1}$. This ends the consideration of the pair $\langle u_0, v_0 \rangle$.

Applying this construction consecutively for all pairs of $u_0 \in P_0$ and $v_0 \in P_1$ (including the pair $\langle 1^n \wedge 0, 0^n \wedge 1 \rangle$) we finally get a system of sets satisfying (1), (3), and (4), which will be denoted still by X_u and P_{uv} .

Step 3. We finally take care of (2). Consider a particular crucial pair $\langle u_0, v_0 \rangle$ in 2^{n+1} . By the density there is a set $P'_{u_0, v_0} \in D_2(n)$, $P'_{u_0, v_0} \subseteq P_{u_0, v_0}$. (In the case when $\langle u_0, v_0 \rangle$ is $\langle 1^n \wedge 0, 0^n \wedge 1 \rangle$ we rather apply Proposition 5 to obtain P'_{u_0, v_0} .)

Define $X'_{u_0} = \text{pr}_1 P'_{u_0, v_0}$ and $X'_{v_0} = \text{pr}_2 P'_{u_0, v_0}$ and spread this change through the chain of crucial pairs in 2^{n+1} . (Note that $X'_{u_0} \equiv X'_{v_0}$ as sets in $\mathbb{P}_{\equiv}^2$ are included in $\equiv$. This keeps $X'_u \equiv X'_v$ for all $u, v \in 2^{n+1}$ through the spreading.)

Executing this step for all crucial pairs in 2^{n+1} we finally end the construction, in $\mathbf{V}^+$, of a system of sets satisfying (1) through (4).

□ (Theorem 3)

References

- [1] L. A. Harrington, A. S. Kechris, A. Louveau. A Glimm – Effros dichotomy for Borel equivalence relations. *J. Amer. Math. Soc.* 1990, 3, pp. 903 – 928.
- [2] L. A. Harrington, D. Marker, S. Shelah, Borel orderings. *Trans. Amer. Math. Soc.* 1988, 310, pp. 293 – 302.